

**READING THIS
MAY SAVE YOUR
ASS**

Shigella is on the rise among
gay/bisexual men

Shigella bacteria are transmitted
by feces - simply put: **SHIT!**

Shigella causes: stomach cramps,
bloody diarrhea, and fever for 4 to 7
days. Got these symptoms??
Call your doctor.

**PROTECT YOURSELF
FROM SHIGELLA**

WASH, WASH, WASH
before, between & after all sex
and "ASS PLAY"

More info: Contact your local
Public Health Department or see the
Health File at:
<http://www.hlth.gov.bc.ca/hlthfile/hfile08n.html>

**READING THIS
MAY SAVE YOUR
ASS**

Shigella is on the rise among
gay/bisexual men

Shigella bacteria are transmitted
by feces - simply put: **SHIT!**

Shigella causes stomach cramps,
bloody diarrhea, and fever for 4 to 7
days. Got these symptoms??
Call your doctor.

**PROTECT YOURSELF
FROM SHIGELLA**

WASH, WASH, WASH
before, between & after all sex
and "ASS PLAY"

More info: Contact your local
Public Health Department or see the
Health File at:
<http://www.hlth.gov.bc.ca/hlthfile/hfile08n.html>

**READING THIS
MAY SAVE YOUR
ASS**

Shigella is on the rise among
gay/bisexual men in British Columbia

Shigella bacteria are transmitted
by feces - simply put: **SHIT!**

Shigella causes stomach cramps,
bloody diarrhea, and fever for 4 to 7
days. Got these symptoms??
Call your doctor.

**PROTECT YOURSELF
FROM SHIGELLA**

WASH, WASH, WASH
before, between & after all sex
and "ASS PLAY"

More info: Contact your local
Public Health Department or see the
Health File at:
<http://www.hlth.gov.bc.ca/hlthfile/hfile08n.html>

READING THIS MAY SAVE YOUR ASS

Shigella Facts

1. Easily transmitted by rimming & even by tiny amounts of shit on:

- penis
- hands
- sex toys
- condoms

2. Who's likely to get sick?

- Anyone, but especially those with weakened immune systems

3. Avoid Shigella

- Wash & use soap
- Question where body parts have been ...touched an ass??

4. Had diarrhea, cramps, fever?

- Contact your doctor
- Inform partners
- Wash hands frequently
- Avoid "sex" until 7 days after symptoms stop
- Avoid handling food that others will eat.

READING THIS MAY SAVE YOUR ASS

Shigella Facts

1. Easily transmitted by rimming & even by tiny amounts of shit on:

- penis
- hands
- sex toys
- condoms

2. Who's likely to get sick?

- Anyone, but especially those with weakened immune systems

3. Avoid Shigella

- Wash & use soap
- Question where body parts have been ...touched an ass??

4. Had diarrhea, cramps, fever?

- Contact your doctor
- Inform partners
- Wash hands frequently
- Avoid "sex" until 7 days after symptoms stop
- Avoid handling food that others will eat.

READING THIS MAY SAVE YOUR ASS

Shigella Facts

1. Easily transmitted by rimming & even by tiny amounts of shit on:

- penis
- hands
- sex toys
- condoms

2. Who's likely to get sick?

- Anyone, but especially those with weakened immune systems

3. Avoid Shigella

- Wash & use soap
- Question where body parts have been ...touched an ass??

4. Had diarrhea, cramps, fever?

- Contact your doctor
- Inform partners
- Wash hands frequently
- Avoid "sex" until 7 days after symptoms stop
- Avoid handling food that others will eat.